

Message

---

**From:** Chism, William [/O=EXCHANGELABS/OU=EXCHANGE ADMINISTRATIVE GROUP (FYDIBOHF23SPDLT)/CN=RECIPIENTS/CN=475879B16C29401A9449DDB69D5F7EB1-WILLIAM CHISM]  
**Sent:** 9/21/2020 4:21:59 PM  
**To:** jnorswor@uark.edu; Stanley Culpepper [stanley@uga.edu]; Bradley, Kevin [bradleyke@missouri.edu]; Orlowski, John [Orlowski.John@epa.gov]; jmcginty@tamu.edu  
**Subject:** Cotton Weed Control Question: dicamba followed by glufosinate

Hi Jason, Joshua, Kevin, and Stanley

We are working on the benefits of dicamba used over the top on dicamba tolerant cotton and we have a what if question. If a user has a heavy population of Palmer amaranth and they have not been controlled after two applications of dicamba what should they try next? Would it be an application of glufosinate or is there another herbicide that might be recommended in you state? We are estimating a weed control budget using dicamba or alternatives and in some areas if they might need an additional trip through the field due to lack of control we were wondering what that might be.

Thank you. Bill and John

---

Bill Chism, Ph.D.  
Senior Biologist  
Biological and Economic Analysis Division  
U.S. EPA/Office of Pesticide Programs  
PII  
[chism.bill@epa.gov](mailto:chism.bill@epa.gov)

John Orlowski, Ph.D.  
Agronomist  
Biological Analysis Branch  
Biological and Economic Analysis Division  
Office of Pesticide Programs  
US Environmental Protection Agency  
206-999-9336